

## On Some General Physical Aspects of the Real Physical Space and Time

A.M. Gevorkian , R. A. Gevorkian\*

*Institute of Physical Research, 378410, Ashtarak-2, Republic of Armenia*

*\*Institute of Radiophysics and Electronics, 378410, Ashtarak-2, Republic of Armenia*

### Abstract

*The paper discusses some aspects of real physical space and time: structure of electrodynamics; structure of gravitational-inert theory in the real world; formation of the real matter; Mach's principle and the radiational cosmic mass; spread and transmission of information in the space of the real world; fundamental and main physical constants, protostar or 3+1-dimensional matter; and some philosophical and general physical approaches to solutions of physical problems.*

1. In the real 4-dimensional space the 3-dimensional and 1-dimensional subspaces are distinguished with the coefficient of imaginary relativity  $ci$  between their dimensions. That is why it is thought of as 3+1-dimensional and relativistic space.

2. Electrodynamics - To the 3-dimensional subspace, determined by 3-dimensional polar vector field (electrical), axial (magnetic) vector is added in the form of a relativistic addition. The union of these vector fields creates the polar-axial (electromagnetic) anti-symmetric tensor of the fourth order. Lagrange's vector equation, which is obtained on the basis of Lagrangian, becomes Maxwell's equations of electrodynamics.

For such a space, as a consequence, the communicational wave equation is obtained which is determined by d'Alembert's operator

$$\square A = \Delta A - \frac{1}{c^2} \frac{\partial}{\partial t^2} A = 0 \quad (1)$$

3. Earlier we showed that an equation is added to Einstein's equation determined by the counter-projection of tensor [1]. Also an expression was received which presented the geometric interpretation of the active principle of equivalence:

$$(-) P_{ik} = P_{ik} \quad (2)$$

Just like the magnetic field is added to the electric field in electrodynamics, here the inert field is added to the gravitational field in the form of equation  $G_{ik} = 2P_{ik}$ . These fields are associated with each other through the principle of active equivalence [1].

4. On the basis of the principle of active equivalence in [1] we received an algebraically transcendental equation, the core of which is illustrated in fig 1.

Fig. 1a presents the equilibrial main state of a virtual particle with real mass equal to zero. On fig. 1b, after deformation of 3+1-dimensional space, another equilibrial state is created, which determines the real, stable matter with a finite mass. The surface of integration is determined from point 1, in the direction of increase of  $x$ , as well as in the direction of decrease of  $1/x$ . The mass obtained this way is the source of gravitational and inert fields, and the electric charge is a necessary attribute during formation of the usual matter [1]. However, it must be noted that the divergence from the real value is approximately 10%, which is due to the spin of the object.

5. In [1] we received and formulated Mach's principle.  $M_i = m_i + \overline{m}_i$  is the effective mass effecting the test body. Relative to the test body, on the leverage  $i$  on both sides, there will always be compensating each other masses  $m_i$  and  $\overline{m}_i$ . Then the effective impact on that leverage is equal to zero, and therefore  $M_i = m_i + \overline{m}_i = 0$ . Naturally, the summation by  $i$  is also equal to zero.

$$\sum M_i = 0 \quad (3)$$

The average value of mass (3) is equal to zero, and the shift from the average value is different from zero. Then  $M = (\sum \overline{\Delta} m_i^2)^{1/2}$  represents the radiational cosmic mass. It creates oscillations in the Metagalactic environment (vacuum oscillations). We think that the real radiation (three-degree equilibrial radiation) is responsible for this phenomenon.

6. The second law of thermodynamics is a typical device for converting information into physical entropy. Information can be viewed as fuel for perpetuum motion of the second type. Virtual particles, which are members of non-excited environment, are in one main state, and the union of these

particles creates macroscopically non-excited vacuum. That is why entropy, which is determined as the logarithm of statistical weight of macroscopic states of subsystems, is equal to zero,  $S = 0$ .

On the basis of Nernst's theorem, we conclude that the non-excited vacuum can be viewed as a thermodynamic system at the absolute zero. For information transmission it is necessary to excite the environment (vacuum), by filling it with excited particles (create fluctuations).

7. The fundamental constants are  $G$ ,  $c$ ,  $h$ . Their combinations can create  $hc$ ,  $c^4/G$ , the fundamental charge and force respectively, formed from the combination of two constants. In [1] the non-measure constant  $\alpha^2 = \frac{hc}{e^2}$  (with 10% accuracy) is received, which associates the real matter with the fundamental constant of space (environment).

Let us consider energy  $e^2/r$  along the entire accessible length (from  $r$  to  $R$ ). It is assumed that the integral represents the fundamental charge, i.e.

$$hc = \int_{r_0}^R \frac{e^2}{r} dr$$

or

$$R = r_0 \exp \frac{hc}{e^2} \quad (4)$$

where  $r_0$  is the fundamental length, , and  $r_0 = \left(\frac{Gh}{c^3}\right)^{1/2}$ ,  $\exp \frac{hc}{e^2} \approx 10^{59 \div 60}$  is the cosmological factor. Then the numerical value of  $R$  is estimated as  $R \sim 10^{27} \text{sm}$ .

The measures of the Metagalaxy are estimated at the same value through astrophysical observations. If the fundamental length in the formula is substituted by the fundamental value of mass, then we receive

$$M = m_0 \exp \frac{hc}{e^2}, \quad M \sim 10^{55} gr \quad (5)$$

The value of the mass of the Metagalaxy is also in accordance with astrophysical estimations.

Inside the atom the integral of Coulomb potential is equal to  $(hc)^{1/2}$ , and is equal to zero outside of the atom.

$$(hc)^{1/2} = \int_{r_{kl}}^R \frac{e}{r} dr \quad \text{or} \quad R = r_{kl} \exp \frac{(hc)^{1/2}}{e}, \quad R \sim 10^{-10} sm$$

The hydrogen atom size is received when  $r_{kl} = \frac{e^2}{mc^2}$  (classical radius of the proton).

Another fundamental constant  $c^4/G$  represents Coulomb's, as well as Newton's forces for fundamental physical values. It is assumed that after formation of the Real World, Newton's force between the Metagalaxy and the hydrogen atom, does not change on the distance equal to the classical radius of the hydrogen atom.

Let us introduce  $R = \frac{e^2}{(m_p m_e)^{1/2} c^2}$  - the value of the classical hydrogen atom radius (analogous to that of charged particles). For Newton's force between the Metagalaxy and the hydrogen atom we receive:

$$F = \frac{GM (m_p m_e)^{3/2} c^4}{e^4} \quad or \quad \frac{c^4}{G} = \frac{GM (m_p m_e)^{3/2} c^4}{e^4}$$

After substituting M with its value received from (5), we obtain:

$$\frac{e^4}{G^2 (m_p m_e)^{3/2}} = \left( \frac{hc}{G} \right)^{1/2} \exp \alpha^2$$

where the dimensionless value:

$$\frac{e^2}{G m_p m_e} = \alpha^{2/3} \exp 2/3 \alpha^2 \sim 10^{-40 \div 41} \quad (6)$$

represents the ratio of Coulomb force to Newton's force in the hydrogen atom.

The numerical results received here surprisingly accurately coincide with experimental data. Thus, main dimensionless physical constants like  $\alpha^2 = \frac{e^2}{hc}$ ,  $\frac{e^2}{G m_p m_e} = \beta^2 = \alpha^{2/3} \exp 2/3 \alpha^2$  and cosmological constants like  $R = r_0 \exp \alpha^2$ ,  $M = m_0 \exp \alpha^2$  - the size and mass of Metagalaxy, and Hubble's constant

$$H = \frac{c}{R} = \frac{c}{r_0 \exp \alpha^2} \quad (7)$$

as well as the radiational width

$$\Delta \nu = \frac{4}{3} \frac{\sigma}{hc} T^4 R \quad (8)$$

are added to fundamental physical constants  $h$ ,  $c$ ,  $G$ , which describe the physical environment. They are also added to such fundamental physical values as

$$r_0 = \left(\frac{Gh}{c}\right)^{1/2}, \quad t_0 = \frac{Gh}{c^5}, \quad l_0 = hc, \quad F_0 = \frac{c^4}{G^5}$$

which are formed by the fundamental physical constants.

According to Stefan-Boltzmann's law, the energy density in each point of the real world is determined by a three-degree equilibrial radiation (relic radiation):

$$E = \frac{4\sigma}{c}T^4 \quad (9)$$

where  $\sigma$  is Stefan-Boltzmann's constant  $\sigma = \frac{\pi^2 k^2}{60\eta^3 c^2}$ .

The energy along the length of the atom (e.g. hydrogen atom) is equal to:

$$E = \frac{4}{3} \frac{\sigma}{c} T^4 R$$

where  $R$  is the dimensionless radius (length) of the atom.

The radiational width of spectral lines in atoms is determined by the constant radiation of the environment

$$E = h\Delta\nu = \frac{4}{3} \frac{\sigma}{c} T^4 R$$

and for the radiational width in specters we have

$$\Delta\nu = \frac{4}{3} \frac{\sigma}{hc} T^4 R \sim 10^8 \quad (10)$$

We also have the cosmological factor

$$A = \exp \alpha^2 \sim 10^{60 \div 61} \quad (11)$$

and the atomic factor

$$a = \exp \alpha = 10^5 \quad (12)$$

8. On the basis of generalization of the theory of non-stationar starts, and the analysis of activity of the nuclei of galaxies, and galaxies with blue excess, allowed V. Ambartsumian to conclude that the process of star formation and non-stationar state of some formations are connected with protostar substance [2], which degrades and becomes usual matter. This process is accompanied by release of enormous amount of energy. There have not been any direct observations of objects with protostar substance.

We think that such protostar substance may be the  $3 + 1$ -dimensional substance, which decomposes into smaller unstable fragments until reaching the state of stable particles (electron, proton, photon), releasing energy. Then these particles form the usual 3-dimensional substance, the most basic of which is the hydrogen atom. Stable configurations, such as white dwarfs, neutron stars (Oppenheimer, Volkov), hyperonic stars (Ambartsumian, Saakian, Vartanian) etc, can not be protostar objects. However, there is one interesting work by Muradian [3,4], which describes the morphology of the macroscopic hadron (two-dimensional heavy hadron), which, due to being unusual (application of nuclear physics laws in astrophysics), may possibly be the protostar substance.

If the proton and the electron are the stable "relic" substance ( $3 + 1$ -dimensional), then in order to understand the protostar macroscopic substance, it is necessary to build the internal electrodynamics of the charged particle. The physical presentation of their microscopic forms (electron and proton), may serve as protostar macroscopic substance (one possibility is the ball lightning). Therefore it is necessary to build the electrodynamics of charged particles, or receive Maxwell's equations (non-traditional), which describe the internal structure of electromagnetic field.

9. a) Let us consider equality

$$P + \overline{P} = 1 \tag{13}$$

The sum of probabilities of a phenomenon and its counter-phenomenon is equal to one.

Example:  $n$  number of shooters have a hypothetical probability of to hit the target  $p_i$ . It is necessary to find the probability of hitting the target in the case of simultaneous shooting. Due to factorization of the counter-probability

$$\overline{P} = \prod_i^n (1 - p_i), \text{ we receive } P = 1 - \prod_i^n (1 - p_i) \quad (14)$$

Problem is solved.

This method allows to solve if not all, then the majority of problems in the theory of probabilities and statistics (for big n).

b) Definition

$$f(\overline{x}) = \overline{f}(x)$$

If a function with a counter-argument becomes its own counter-function, such a function is called a counter-odd function.

$$f(\overline{x}) + \overline{f}(x) = 1$$

One of the particular solutions of the functional equation is

$$f(x) = \sin^2 x, \quad x + \bar{x} = \frac{\pi}{2}, \quad x + \bar{x} = 1$$

The simultaneous happening of the phenomenon and the counter-phenomenon takes place in anti-phase. If we imagine the electric and magnetic fields to be the phenomenon and the counter-phenomenon respectively, then Maxwell's electrodynamics can be presented through the model of the example.

c) In [1] we received

$$G_{ik} + \overline{G}_{ik} = 2P_{ik}$$

where  $P_{ik}$  is the tensor of counter-projection, and for scalars the tensor equation becomes equation (14).

$$G + \overline{G} = 1$$

We received that the sum of scalar curvature  $G$  and curvature  $\overline{G}$  is equal to one. In the framework of this model we received the equation of the inert field.

Cases a) and c) can be brought to case b) if we assume that the object and the counter-object change and depend on the parameter of state. More

examples can be brought which are described by this model. Based on these examples, the general law can be formulated.

The core of the law of dialectics on "struggle and unity of opposites" lies not so much in the "struggle", rather it lies in their unity. This means that in the process of the "struggle", one opposition becomes enriched at the account of the other one. Thus, oppositions struggle, and in this process they turn into each other either partially or completely. These changes happen in the anti-phase. The process of opposing may be stationar, like in examples a) and c), or non-stationar, as in b). If we assume that objects described in examples a) and c) are non-stationar, then they also can be brought to example b). Electrodynamics is described by the model of example b), where the object is the electric field, and the counter-object is the magnetic field. Maxwell's equations form a unified system where the electric and the magnetic fields oppose to each other and change in the anti-phase. As a consequence of the unified system the wave equation is received.

$$\square A = 0 \quad (15)$$

If we assume that in examples a) and c) the objects are non-stationar, then for the first example the wave equation of amplitudes of probabilities is expected, while for the third example the wave equation of the symmetric tensor  $g_{ik}$  is expected.

$$\square g_{ik} = 0 \quad (16)$$

Similarly to equations for 4-dimensional vector potential, the electromagnetic field can also be generalized into a 10-component tensor equation like (16). Rules of d'Alembertian effect on tensor are not yet completely clear. It is necessary that construction  $\square$  be some kind of general differential operator of the second order, which contains a member like

$$\frac{\partial^2 g_{ik}}{\partial x^\mu \partial x^\nu} = 0$$

This operator requires further mathematical work. In [1] we used extreme values of the object and the counter-object and received two pairs of equivalent equations:

$$G_{ik} + 2 \frac{(g_{ik} - \delta_{ik})}{\xi_i \xi^i} = 0, \quad \overline{G}_{ik} = 0, \quad \text{First equivalent pair}$$



$$\overline{G}_{ik} - 2 \frac{(g_{ik} - \delta_{ik})}{\xi_i \xi^i} = 0, \quad G_{ik} = 0, \quad \textit{Second equivalent pair}$$

which describes gravitational and inert fields.

Examples used in point 9 demonstrate that the object and the counter-object coexist and form a unified system. The real physical space and time are described by such general physical aspects.

#### Literature

1. A. M. Gevorkian, R.A. Gevorkian. "Inertia in the Structure of Four-dimensional Space". Automated e-print archives at arXiv.org: Physics, General Relativity and Quantum Cosmology, gr-qc/0002046, 14.2.2000 xxx.lanl.gov.
2. V. A. Ambarcumyan "About Protostars" Reports of Academy of Science (DAN), 16, 97, 1953
3. R. M. Muradian "Astrophysics" 11, 237, 1975
4. R. M. Muradian "Astrophysics" 13, 63 1975

This figure "Gevorkian3.png" is available in "png" format from:

<http://arXiv.org/ps/physics/0101084v1>